

INF4140, 2012. Exercise 7

While program

Consider the following program S :

```
S: x = 0; y = b;
   while (x < y) { x = x + 2; y = y + 1 }
```

Prove that the following triple is a theorem in PL:

$$\{b \geq 0\} S \{x == 2 * b\}$$

You may use the following predicate I as loop invariant:

$$I : x \leq y \wedge x == 2(y - b)$$

Factorial function

Consider the following program S :

```
S: i = 0; x = 1;
   while (i < n) {
     i = i + 1;
     x = x * i;
   }
```

Prove the following triple using PL:

$$\{n \geq 0\} S \{x == n!\}$$

As a loop invariant I , you may use: $I : x == i! \wedge i \leq n$.

You may assume the following when reasoning about the factorial function:

- 1) $0! == 1$
- 2) $(j + 1)! == j! * (j + 1)$ for any integer $j \geq 0$

Monitor verification

Consider the monitor for Shortest-Job-Next allocation in the book (section 5.2.3). Use Programming Logic, extended with rules for **signal** and **wait** (lecture slides, week 6), to prove that this monitor satisfies the second part of the SJN invariant:

$$\mathbf{free} \Rightarrow (\#\mathbf{turn} == 0)$$

(You may use the rule for **wait(cv)** to reason about **wait(cv,rank)**).

Hint. When arriving at an implication, it is enough to argue for the truth of it. However, we may use the following rules when reasoning about implications.

$$\frac{}{\mathbf{false} \Rightarrow A} \qquad \frac{(A \wedge B) \Rightarrow C}{A \Rightarrow (B \Rightarrow C)} \qquad \frac{(A \wedge B) \Rightarrow C}{((A \Rightarrow B) \wedge A) \Rightarrow C} \qquad \frac{(\neg A) \vee B}{A \Rightarrow B}$$

From the book

2.22, 2.16, 2.24, 2.31, (2.28a, 2.29a)