

- 2 -

SOLUTIONS MAT 2700 MID-TERM ASSIGNMENT  
October 2008

Exercise 1a)

In general we have

$$R_n = \frac{S_n(1) - S_n(0)}{S_n(0)}$$

Here this gives

$$R_1(\omega_1) = \frac{4-3}{3} = \underline{\underline{\frac{1}{3}}}, \quad R_1(\omega_2) = \frac{2-3}{3} = \underline{\underline{-\frac{1}{3}}}$$

Exercise 1b A probability measure  $Q$  is risk neutral iff

$$E_Q \left[ \frac{R_{n+r}}{1+r} \right] = 0 \quad \text{for all } n = 1, \dots, N$$

Here this gives the equations ( $Q = (Q_1, Q_2)$ )

$$\begin{cases} Q_1 + Q_2 = 1 \\ \frac{1}{3} Q_1 - \frac{1}{3} Q_2 = \frac{1}{10} \end{cases}$$

This system has the unique solution

$$\underline{\underline{Q_1 = Q(\omega_1) = \frac{13}{20}}}, \quad \underline{\underline{Q_2 = Q(\omega_2) = \frac{7}{20}}}$$

Exercise 1c Since there exists a risk neutral probability measure  $Q$ , there is no arbitrage.

Exercise 1d) Since there are no arbitrages, there cannot be any dominant strategies.

Exercise 1e) Since there is only one risk neutral probability measure, the market is complete.

Exercise 1f) Since there exists only one risk neutral probability measure, the market satisfy the law of one price.

Exercise 2a In the market  $M_k$  the returns are

$$R_1^k(w_1) = \frac{1}{3} + k, \quad R_1^k(w_2) = -\frac{1}{3} + k$$

Therefore the equations for a risk neutral probability measure  $Q^{(k)} = (q_1, q_2)$  are

$$\begin{cases} q_1 + q_2 = 1 \\ (\frac{1}{3} + k)q_1 + (-\frac{1}{3} + k)q_2 = \frac{1}{10} \end{cases}$$

The solution of this system is

$$q_1 = Q^{(k)}(w_1) = \frac{13}{20} - \frac{3}{2}k, \quad q_2 = Q^{(k)}(w_2) = \frac{7}{20} + \frac{3}{2}k,$$

where

$$\underline{\underline{0 \leq k < \frac{13}{30}}}$$

Exercise 2b)

(i) If  $K < \frac{13}{30}$  there exists a risk neutral probability measure and hence there is no arbitrage

(ii) If  $K \geq \frac{13}{30}$  then no risk neutral probability measures exist. Hence the market has an arbitrage.

Exercise 2c)

(i) If  $K < \frac{13}{30}$  there exists a unique risk neutral probability measure and hence the market is complete.



Exercise 2d)

Maximize  $E[\ln W]$  over all  $W \in \mathbb{R}^2$ ,  
 subject to  $E_{Q^{(k)}} \left[ \frac{W}{B_1} \right] = v$

Using the Lagrange multiplier method we first maximize

$$h(w) := E[\ln W] - \lambda E_{Q^{(k)}} \left[ \frac{W}{B_1} \right]$$

without constraints.

$$\text{Now } h(W) = E \left[ \ln W - \lambda \frac{Q^{(k)}}{P} \frac{W}{B_1} \right]$$

We can maximize

$$g(x) = \ln x - \lambda \frac{Q^{(k)}(w)}{P(w)} \frac{x}{B_1(w)}$$

for each  $w$ .

The first order condition is

$$0 = g'(x) = \frac{1}{x} - \lambda \frac{L^{(k)}(w)}{B_1(w)} \quad ; \quad L^{(k)}(w) = \frac{Q^{(k)}(w)}{P(w)}$$

which gives

$$x = \hat{W}^{(k)} = \frac{B_1(w)}{\lambda L^{(k)}(w)}$$

The constraint

$$E \left[ L^{(k)} \frac{\hat{W}^{(k)}}{B_1} \right] = v$$

gives the value of  $\lambda$ :

$$E \left[ \frac{L^{(k)} B_1}{\lambda L^{(k)} B_1} \right] = v \Rightarrow \lambda = \frac{1}{v} =: \hat{\lambda} =$$

Hence the optimal  $W = \hat{W}(\hat{x})$  is given by

$$\hat{W}(\hat{x}) = v \frac{B_1}{L^{(k)}} \quad (\text{as in Exercise 2.2 in [P]})$$

With the given values for  $P$ ,  $B_1$ , and  $Q^{(k)}$  this becomes

$$\hat{W}(\hat{x})(w_1) = v \frac{\frac{1}{2} \cdot (1 + \frac{1}{10})}{\frac{13}{20} - \frac{3}{2}k} = \frac{11v}{13 - 30k}$$

Similarly,

$$\hat{W}(\hat{x})(w_2) = v \frac{\frac{1}{2} (1 + \frac{1}{10})}{\frac{7}{20} + \frac{3}{2}k} = \frac{11v}{7 + 30k}$$

$$0 \leq k < \frac{13}{30}$$

Exercise 2e) We want to find  $F^{(k)}$  such

that

$$v(1+r + F^{(k)}(R_1^{(k)} - r)) = \hat{W}(\hat{x})$$

i.e.,

$$(i) \quad v(1 + \frac{1}{10} + F^{(k)}(\frac{1}{3} + k - \frac{1}{10})) = \frac{11v}{13 - 30k} \quad (w = w_1)$$

and

$$(ii) \quad v(1 + \frac{1}{10} + F^{(k)}(-\frac{1}{3} + k - \frac{1}{10})) = \frac{11v}{7 + 30k} \quad (w = w_2)$$



Only one of these two equations is necessary to determine  $F^{(k)}$ , Equation (i) gives

$$\frac{11}{10} + F^{(k)} \left( \frac{7}{30} + k \right) = \frac{11}{13 - 30k}$$

or

$$F^{(k)} = \frac{99(-1 + 10k)}{(7 + 30k)(13 - 30k)}$$

The corresponding optimal utility is

$$J_k^{(v)} = E[\ln \hat{W}^{(k)}] =$$

$$= \ln v + \ln 11 - \frac{1}{2} \ln(13 - 30k) - \frac{1}{2} \ln(7 + 30k)$$

Exercise 3 We now want to solve the constrained maximum problem

$$\text{maximize } E[\ln(V_1^{(F)})]$$

subject to

$$F \geq 0 \quad \text{and} \quad V_0 = v$$

To this end, we introduce the markets  $\mathcal{M}_k$  in Exercise and we solve the unconstrained problem

$$\left\{ \begin{array}{l} \text{maximize } E[\ln V_1^{(F)}] \\ \text{(given that } V_0 = v) \end{array} \right.$$

in the market  $\mathcal{M}_k$ .

- \* -

This was done in Exercise 2.

We now seek  $k = \hat{k}$  such that the corresponding optimal portfolio  $F(\hat{k})$  satisfies the conditions

$$(i) \quad F(\hat{k}) \geq 0$$

and

$$(ii) \quad F(\hat{k}) \cdot \hat{k} = 0$$

(Here  $d(k) = \sup_{F \geq 0} (-F) \cdot k = 0$  for all  $k \in \tilde{K} = [0, \infty)$ )

Equation (ii) tells us that either  $F(\hat{k}) = 0$  or  $\hat{k} = 0$ .

The case  $\hat{k} = 0$  is impossible, because it makes  $F(\hat{k}) < 0$ . Hence we must choose  $\hat{k}$  such that

$$\underline{F(\hat{k}) = 0} \quad \text{i. e.} \quad \underline{\hat{k} = \frac{1}{10}}$$

Hence  $F(\hat{k}) = 0$  is the optimal portfolio for the constrained maximum problem in Exercise 3. The corresponding optimal utility is

$$J_{\tilde{K}}(v) = E[\ln(v(1+r))] = \ln v + \ln \frac{11}{10}.$$



Alternatively, one could find  $\hat{k}$  by minimizing  $J_k(v)$  over all  $k \in [0, \frac{13}{30})$ :

Define

$$h(k) = \ln(13 - 30k) + \ln(7 + 30k)$$

Then

$$h'(k) = \frac{-30}{13 - 30k} + \frac{30}{7 + 30k} = 0$$

iff

$$-(7 + 30k) + 13 - 30k = 0$$

i.e.

$$\underline{\underline{k = \hat{k} = \frac{1}{10}}}$$

Exercise 4 a) To find all risk neutral probability measures  $Q = (q_1, q_2, q_3)$  we solve the system of equations

$$\begin{aligned} q_1 + q_2 + q_3 &= 1 && (Q \text{ prob. meas.}) \\ \frac{1}{3} q_1 + (-\frac{1}{3}) q_2 + 0 \cdot q_3 &= 0 && (E_Q[R] = 0) \end{aligned}$$

which has the solution

$$\begin{cases} q_1 = Q(\omega_1) = \frac{1}{2} - \frac{1}{2} q_3 \\ q_2 = Q(\omega_2) = \frac{1}{2} - \frac{1}{2} q_3 \\ q_3 = Q(\omega_3) \in (0, 1) \end{cases}$$



Exercise 4b) A claim  $X = (X_1, X_2, X_3)$   
is attainable iff

$$E_Q[X]$$

has the same value for all risk neutral probability measures  $Q$ .

With  $Q$  as in 4a) we get

$$\begin{aligned} E_Q[X] &= \left(\frac{1}{2} - \frac{1}{2}q_3\right)X_1 + \left(\frac{1}{2} - \frac{1}{2}q_3\right)X_2 + q_3X_3 \\ &= \frac{1}{2}X_1 + \frac{1}{2}X_2 + q_3\left(-\frac{1}{2}X_1 - \frac{1}{2}X_2 + X_3\right) \end{aligned}$$

This is independent of  $q_3$  iff

$$-\frac{1}{2}X_1 - \frac{1}{2}X_2 + X_3 = 0$$

or

$$\underline{\underline{X_1 + X_2 - 2X_3 = 0}}$$

Exercise 4c) We seek  $(H_0, H_1)$  such that

$$H_0(1+r) + H_1S_1(1, \omega) = X(\omega) =$$

i.e.

$$\left\{ \begin{array}{l} H_0 + 4H_1 = 3 \\ H_0 + 2H_1 = 1 \\ H_0 + 3H_1 = 2 \end{array} \right.$$

which has the solution  $\underline{\underline{H_0 = -1}}, \underline{\underline{H_1 = 1}}$ .

