

FYS 4110 Modern Quantum Mechanics, Fall Semester 2016

Problem set 1

We begin the weekly sets with some problems concerning basic and useful mathematical relations.

1.1 Commutators and anti-commutators

We use the standard notation for commutators and anticommutators

$$[A, B] = AB - BA \quad \{A, B\} = AB + BA \quad (1)$$

where A and B are two operators or matrices. Show the following identities,

$$\begin{aligned} [A, BC] &= [A, B]C + B[A, C] \\ \{A, BC\} &= \{A, B\}C - B\{A, C\} \end{aligned} \quad (2)$$

1.2 Trace and determinant

We remind you about the following relations

$$\text{Tr}(AB) = \text{Tr}(BA), \quad \det(AB) = \det A \det B \quad (3)$$

a) Assume \hat{A} to be a quantum observable and A to be the matrix representation of the observable in an orthonormalized basis $\{|n\rangle\}$, which means

$$A_{mn} = \langle m | \hat{A} | n \rangle \quad (4)$$

We define the trace and determinant of the (abstract) operator as

$$\text{Tr } \hat{A} = \text{Tr } A, \quad \det \hat{A} = \det A \quad (5)$$

Show that if we change to a new basis $\{|n'\rangle\}$, which is related to the first by a unitary transformation, that will not change the values of the trace and determinant.

b) Assume \hat{A} is a hermitian operator with eigenvalues $a_n, n = 1, 2, \dots$. Explain why the trace and determinant can be expressed in terms of the eigenvalues as

$$\text{Tr } \hat{A} = \sum_n a_n \quad \det \hat{A} = \prod_n a_n \quad (6)$$

c) The *spectral decomposition* of an hermitian operator \hat{A} is a sum of the form

$$\hat{A} = \sum_n a_n |n\rangle \langle n| \quad (7)$$

where a_n are the eigenvalues and $|n\rangle$ are the corresponding eigenvectors of the operator. A function $f(a)$ defines an *operator function* $\hat{f} \equiv f(\hat{A})$ of \hat{A} by the related decomposition

$$\hat{f} \equiv \sum_n f(a_n) |n\rangle \langle n| \quad (8)$$

Use this definition and the results of problem b) to show that we have the following relation

$$\det e^{\hat{A}} = e^{\text{Tr } \hat{A}} \quad (9)$$

We assume the trace of \hat{A} to be well defined and finite (which may not necessarily be the case in an infinite dimensional Hilbert space).

d) Show that for general state vectors $|\psi\rangle$ and $|\phi\rangle$ we have the relation

$$\langle\psi|\phi\rangle = \text{Tr}(|\phi\rangle\langle\psi|) \quad (10)$$

1.3 Dirac's delta function

The basic relation defining the delta functions is the following

$$f(x) = \int_{-\infty}^{\infty} dx' \delta(x - x') f(x') \quad (11)$$

with $f(x)$ as any chosen function. Clearly $\delta(x)$ is not a function in the usual sense, and in particular it has the property that $\delta(x) = 0$ for $x \neq 0$ and $\delta(0) = \infty$. Nevertheless it is possible (with some care) to treat it as a function, and as we know from the wavefunction description of quantum physics it is in many cases a very useful concept.

a) We remind you about the formulas for Fourier transformation in one dimension

$$f(x) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} dk \tilde{f}(k) e^{ikx} \quad (12)$$

$$\tilde{f}(k) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} dx f(x) e^{-ikx} \quad (13)$$

Show that the delta function has the following Fourier expansion

$$\delta(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} dk e^{ikx} \quad (14)$$

b) Assume $g(x)$ is a differentiable function with zero at one point x_0 ,

$$g(x_0) = 0 \quad \text{for} \quad (15)$$

Assume also that the derivative does not vanish at this point, $g'(x_0) \neq 0$. Show by use of the definition (11), and by studying the integral $\int dx \delta(g(x)) f(x)$, that we have the following relation

$$\delta(g(x)) = \frac{1}{|g'(x_0)|} \delta(x - x_0) \quad (16)$$

(Hint, make change of variable $x \rightarrow g$ in the integral.) Assume that the function $g(x)$ has several zeros, at the points $x = x_i$. Explain why this gives the generalized formula

$$\delta(g(x)) = \sum_i \frac{1}{|g'(x_i)|} \delta(x - x_i) \quad (17)$$

1.4 Position and momentum eigenstates

The position and momentum eigenstates are given by the relations

$$\hat{x}|x\rangle = x|x\rangle \quad \langle x|x'\rangle = \delta(x - x') \quad \int dx |x\rangle\langle x| = \mathbb{1} \quad (18)$$

$$\hat{p}|p\rangle = p|p\rangle \quad \langle p|p'\rangle = \delta(p - p') \quad \int dp |p\rangle\langle p| = \mathbb{1} \quad (19)$$

Furthermore, in the x -representation the momentum operator is given by $\hat{p} = -i\hbar \frac{d}{dx}$. Use these relations together with the Fourier expansion of the delta function to show that the scalar product of a momentum and a position state is given by

$$\langle x|p\rangle = \frac{1}{\sqrt{2\pi\hbar}} e^{\frac{i}{\hbar}xp} \quad (20)$$