

FYS3520 - Problem set 1

Spring term 2017

Problem 1 – Discussion in class

a) Derive following commutator relations. What is their significance?


$$\begin{aligned}[\widehat{L}_x, \widehat{L}_y] &= i\hbar\widehat{L}_z, & [\widehat{L}_x, \widehat{y}] &= i\hbar\widehat{z}, & [\widehat{L}_x, \widehat{p}_y] &= i\hbar\widehat{p}_z, \\ [\widehat{L}_x, \widehat{x}] &= [\widehat{L}_x, \widehat{p}_x] = [\widehat{L}_x, \widehat{L}^2] = [\widehat{L}_x, \widehat{r}^2] = [\widehat{L}_x, \widehat{p}^2] &= 0\end{aligned}$$

Hint: You may use $\mathbf{L} = \mathbf{r} \times \mathbf{p} \rightarrow \widehat{\mathbf{L}} = \widehat{\mathbf{r}} \times \widehat{\mathbf{p}}$ and $[\widehat{r}_i, \widehat{p}_j] = i\hbar\delta_{ij}$

b) Given a system with the angular momentum $L = 0$ ($L = 2$) and spin $S = 1/2$, write down the states and their degeneracy in spectroscopic notation: $^{2S+1}L_J$.

Problem 2 – Discussion in class


We want to get a qualitative understanding of the situation where an incoming free particle from the left with energy E hits on a potential barrier $V(x)_{II} > 0$. The particle can be taken as wave-package produced by a superposition of plain waves. In the following graphs, two such cases are plotted for a the real part of the function, $Re(\psi(x)) = \cos(k_I x)$. Sketch how the wave will continue in the different regions and compare the situation to a classical system.



Problem 3 – Discussion in class

In a simple model, the nuclear potential can be assumed to be an **infinite spherical well**. The solutions of the Schrödinger equation are then given by product of the radial function $R(kr)$, which are proportional to the Bessel functions $j_l(kr)$, and the spherical harmonics $Y_l^m(\vartheta, \varphi)$.

- What is the meaning of the different symbols? What quantum numbers does the energy of the states depend on? What is there degeneracy?
- What is the relation between the quantum numbers?
- Sketch the probability density $|\psi_{n,l,m}|^2$ for i) $n=0, l=0, m=0$ and ii) $n=2, l=1, m=1$.
- What is the parity of these states?



Problem 4

Use the notation ${}^A_Z X_N$ with mass number A , element number Z and neutron number N .

- How many neutrons does ${}^{232}\text{Th}$ and ${}^{235}\text{U}$ have?
- List all **stable** nuclei with mass number 20.
- List all **stable** nuclei with element number 82.
- List all **stable** polonium (Po) nuclei.
- Write down all the **stable** Zirconium (Zr) isotopes.

Hint: Charts of most known nuclides can be found at <http://www.nndc.bnl.gov/chart/chartNuc.jsp> or at <https://www-nds.iaea.org/livechart>.

Problem 5

Of the following nuclei: ${}^{16}\text{O}$, ${}^{19}\text{Ne}$, ${}^{18}\text{F}$, ${}^{15}\text{N}$, ${}^{16}\text{C}$, ${}^{20}\text{O}$, ${}^{16}\text{Ne}$, ${}^{14}\text{C}$, ${}^{20}\text{N}$ and ${}^{20}\text{Ne}$.

- Write out the full ${}^A_Z X_N$ notation for all the listed nuclei.
- List the nuclei that are **isotopes**.
- List the nuclei that are **isotones**.
- List the nuclei that are **isobars**.
- What is an **isomer**?

Problem 6

What is the mass of one ${}^{12}\text{C}$ and ${}^{239}\text{Pu}$ nucleus in

- Atomic mass units (AMU) [u]?
- Units of $[\text{MeV}/c^2]$?
- In [kg]?

Hint: There are several databases containing properties of known nuclei. The Atomic Mass Data Center (AMDC) have lists of masses of most of the known nuclides available as a text file at <https://www-nds.iaea.org/amdc/> or as an interactive table at <https://www-nds.iaea.org/livechart>.

Problem 7

The **one-dimensional** time-independent Schrödinger equation is given by

$$-\frac{\hbar^2}{2m} \frac{d^2\psi(x)}{dx^2} + V(x)\psi(x) = E\psi(x)$$

- Derive this from the time dependent Schrödinger equation for a time-independent potential $V(x, t) = V(x)$.
- Assume that the potential is a infinite well, ie. infinite at $x < 0$ and $x > a$ and zero in between ($0 < x < a$). Show that the eigenstates are given by

$$\psi_n = \sqrt{\frac{2}{a}} \sin \frac{n\pi x}{a}, \quad n = 1, 2, 3, \dots$$

and find the energy E_n for each state.

- Assume that we have a particle in the super-position: $\Psi(x) = A(\psi_1(x) + \psi_3(x))$. Find the normalization factor A .
- Find the average energy of $\Psi(x)$. What is the variance?
- What is the probability of finding the particle between $\frac{a}{4} \leq x \leq \frac{3a}{4}$?

Problem 8 – A nice little extra

The potential of a simple one-dimensional harmonic oscillator in one dimension can be described by $V(x) = \frac{1}{2}m\omega^2x^2$.

- Find the energy of the number state $|n\rangle$ using ladder operators:

$$\hat{a} = \frac{1}{\sqrt{2\hbar m\omega}}(m\omega\hat{x} + i\hat{p})$$
$$\hat{a}^\dagger = \frac{1}{\sqrt{2\hbar m\omega}}(m\omega\hat{x} - i\hat{p})$$

Remember: $\hat{a}|n\rangle = \sqrt{n}|n-1\rangle$, $\hat{a}^\dagger|n\rangle = \sqrt{n+1}|n+1\rangle$, and $[\hat{a}, \hat{a}^\dagger] = 1$.

- What two particle states with fermions are possible with the three first states (ignoring spin)?
- Give the degeneracy for 3D symmetric HO, so $\omega_x = \omega_y = \omega_z$, with $E = \frac{7}{2}\hbar\omega$.