

## FYS 3610

Exercise Week 34 due 31. August 2016

## Exercises

In a static magnetic field  $\vec{B}_0$  the equation of motion of a single charged particle is given by:

$$m\frac{d\vec{v}}{dt} = q(\vec{v}\times\vec{B}_0).$$

The position vector  $\vec{r}$  is then found from

$$\frac{d\vec{r}}{dt} = \vec{v}.$$

To study the motion of the particle in the plane perpendicular to the magnetic field, we will assume that is oriented along the *z*-axis, i.e.,  $\vec{B}_0 = (0,0,B_0)$ . The equation of motion then simplifies to

$$\frac{dv_x}{dt} = \frac{q}{m}v_y B_0$$
$$\frac{dv_y}{dt} = -\frac{q}{m}v_x B_0$$

 $\frac{dr_x}{dt} = v_x$ 

 $\frac{dr_y}{dt} = v_y.$ 

and

To solve these equations numerically we use Euler's method to approximate the time derivative as the difference of the function value between time t and the next time step t + h, where h is the step size:



$$\frac{df(t)}{dt} \approx \frac{f(t+h) - f(t)}{h}$$

Applying this to the equations of motion gives

$$\frac{v_x(t+h) - v_x(t)}{h} = \frac{q}{m} v_y(t) B_0$$
$$\frac{v_y(t+h) - v_y(t)}{h} = -\frac{q}{m} v_x(t) B_0$$

and

$$\frac{r_x(t+h) - r_x(t)}{h} = v_x$$
$$\frac{r_y(t+h) - r_y(t)}{h} = v_y.$$

We can rearrange these equations such that we find an expression for  $v_x$ ,  $v_y$ ,  $r_x$ , and  $r_y$  at the next time step, i.e., at t + h, depending on the function values from the previous time step t:

$$v_x(t+h) = v_x(t) + h\left(\frac{q}{m}v_y(t)B_0\right)$$
$$v_y(t+h) = v_y(t) - h\left(\frac{q}{m}v_x(t)B_0\right)$$
$$r_x(t+h) = r_x(t) + hv_x(t)$$
$$r_y(t+h) = r_y(t) + hv_y(t).$$

**Exercise 1:** Write a computer program in the language of your choice (preferably python) that numerically solves the equation of motion for an electron and an oxygen ion (O<sup>+</sup>) in a static magnetic field of 50,000 nT. Let the initial position vector be  $\vec{r} = (0,0,0)$  and the initial velocity vector  $\vec{v} = (500,0,0)$  m/s, for both particles. Choose your time step *h* such that the gyro motion is properly resolved!

**Exercise 2:** Plot the particle's trajectories and check the theoretical predictions of the gyro radius and gyro frequency!