

UNIVERSITY OF OSLO

Faculty of Mathematics and Natural Sciences

Exam in INF5830 - Natural language processing

Day of exam: 15 June 2015

Exam hours: at 14:30 – 4 hours

This examination paper consists of 3 pages including this.

Appendices:

Statistical formulas – 1 page

Statistical table – 4 pages


Permitted materials: None

*Make sure that your copy of this examination paper
is complete before answering.*

- You may answer in English, Norwegian, Danish or Swedish.
- You should answer all questions. The weight of the various questions are indicated.
- You should read through the whole set to see whether anything is unclear so that you can ask your questions to the teachers when they arrive.
- If you think some assumptions are missing, make your own and explain them!

1 Dependency syntax and parsing (25 %)

- (a) Describe the formal conditions on well-formed dependency graphs.
- (b) Consider the dependency graph for the English sentence *He heard a song today that he disliked*:



Is the graph above projective? Why or why not?

- (c) What are the main differences between graph-based and transition-based approaches to data-driven dependency parsing? What type of parser is Maltparser?

2 Semantic role labeling (25 %)

- (a) Provide three roles from the traditional set of semantic roles along with a short description. Apply these to the sentence from exercise 1.
- (b) Consider the following frame taken from FrameNet:

Definition: An Agent cuts an Item into Pieces using an Instrument:	
Frame Elements (core):	
Agent	person cutting the Item
Item	item being cut
Pieces	parts of the original Item
Frame Elements (non-core):	
Instrument	instrument with which the Item is cut
Manner	manner in which Item is cut
Place	where cutting takes place
Purpose	purpose for cutting
Lexical Units:	<i>carve, chop, cube, cut, dice fillet, mince, pare, slice</i>

- (i) Briefly describe the notion of a *frame* and the different frame components above.

- (ii) In what way is Fillmore's frame semantics a response to criticism of the traditional set of semantic roles?
- (c) In Semantic Role Labeling (SRL), parsing is often used as a pre-processing step for feature construction. Provide two examples of features commonly used in SRL that require syntactic parsing and use the dependency graph above to give one example for each of the features you describe.

3 Estimation (15%)

Kim has constructed a classifier and is testing it on a test set with 400 items. It achieves an accuracy of 0.64. Estimate a confidence interval for the accuracy of the classifier at level 0.99.

(A little help since you do not have a computer at hand: $\sqrt{0.64 \times 0.36} = 0.48$)

4 Collocations (20%)

Explain what a collocation is. There have been proposed several different measures for the association between the two words in a collocation. Describe two or three of them.

5 Machine learning (15%)

Multinomial logistic regression is a method for classification which is also called Maximum entropy classification. It is similar to Naive Bayes classification and can in some respects be considered a refinement of Naive Bayes classification. When trained on the same training material with the same set of features, the Logistic regression classifier should in principle perform at least as well as the Naive Bayes classifier when measured on the training material, and sometimes it will perform better. Without going deeply into the mathematical details, explain why this is so.

Also, explain why adding more features may deteriorate the results of the Naive Bayes classifier, but not of the Logistic regression classifier.

In spite of this, Jurafsky and Martin claim "Furthermore, naive Bayes works extremely well (even better than logistic regression or SVMs) on small datasets or short documents." Why do you think that is the case?

END

INF5830, 2015, some statistical formulas

Z-score

Given a normal distribution with mean μ and standard deviation σ . The Z-score of a data point x

$$Z = \frac{x - \mu}{\sigma}$$

expresses the distance of x from μ in terms of standard deviations.

t-test

The t-statistics

$$t = \frac{\bar{x} - \mu}{\sqrt{\frac{s^2}{n}}}$$

where

- \bar{x} is the mean of a simple random sample
- n is the size of the sample
- s is the sample standard deviation

Two sample t-test

$$t = \frac{\bar{x}_1 - \bar{x}_2}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}$$

Standard deviation of proportion

When p is a proportion $\frac{k}{n}$ (k successes out of n), the variance is

$$p(1 - p)$$

STATISTICAL TABLES

Cumulative normal distribution

Critical values of the t distribution


Critical values of the F distribution

Critical values of the chi-squared distribution

TABLE A.1

Cumulative Standardized Normal Distribution

$A(z)$ is the integral of the standardized normal distribution from $-\infty$ to z (in other words, the area under the curve to the left of z). It gives the probability of a normal random variable not being more than z standard deviations above its mean. Values of z of particular importance:



z	$A(z)$	
1.645	0.9500	Lower limit of right 5% tail
1.960	0.9750	Lower limit of right 2.5% tail
2.326	0.9900	Lower limit of right 1% tail
2.576	0.9950	Lower limit of right 0.5% tail
3.090	0.9990	Lower limit of right 0.1% tail
3.291	0.9995	Lower limit of right 0.05% tail

z	0.00	0.01	0.02	0.03	0.04	0.05	0.06	0.07	0.08	0.09
0.0	0.5000	0.5040	0.5080	0.5120	0.5160	0.5199	0.5239	0.5279	0.5319	0.5359
0.1	0.5398	0.5438	0.5478	0.5517	0.5557	0.5596	0.5636	0.5675	0.5714	0.5753
0.2	0.5793	0.5832	0.5871	0.5910	0.5948	0.5987	0.6026	0.6064	0.6103	0.6141
0.3	0.6179	0.6217	0.6255	0.6293	0.6331	0.6368	0.6406	0.6443	0.6480	0.6517
0.4	0.6554	0.6591	0.6628	0.6664	0.6700	0.6736	0.6772	0.6808	0.6844	0.6879
0.5	0.6915	0.6950	0.6985	0.7019	0.7054	0.7088	0.7123	0.7157	0.7190	0.7224
0.6	0.7257	0.7291	0.7324	0.7357	0.7389	0.7422	0.7454	0.7486	0.7517	0.7549
0.7	0.7580	0.7611	0.7642	0.7673	0.7704	0.7734	0.7764	0.7794	0.7823	0.7852
0.8	0.7881	0.7910	0.7939	0.7967	0.7995	0.8023	0.8051	0.8078	0.8106	0.8133
0.9	0.8159	0.8186	0.8212	0.8238	0.8264	0.8289	0.8315	0.8340	0.8365	0.8389
1.0	0.8413	0.8438	0.8461	0.8485	0.8508	0.8531	0.8554	0.8577	0.8599	0.8621
1.1	0.8643	0.8665	0.8686	0.8708	0.8729	0.8749	0.8770	0.8790	0.8810	0.8830
1.2	0.8849	0.8869	0.8888	0.8907	0.8925	0.8944	0.8962	0.8980	0.8997	0.9015
1.3	0.9032	0.9049	0.9066	0.9082	0.9099	0.9115	0.9131	0.9147	0.9162	0.9177
1.4	0.9192	0.9207	0.9222	0.9236	0.9251	0.9265	0.9279	0.9292	0.9306	0.9319
1.5	0.9332	0.9345	0.9357	0.9370	0.9382	0.9394	0.9406	0.9418	0.9429	0.9441
1.6	0.9452	0.9463	0.9474	0.9484	0.9495	0.9505	0.9515	0.9525	0.9535	0.9545
1.7	0.9554	0.9564	0.9573	0.9582	0.9591	0.9599	0.9608	0.9616	0.9625	0.9633
1.8	0.9641	0.9649	0.9656	0.9664	0.9671	0.9678	0.9686	0.9693	0.9699	0.9706
1.9	0.9713	0.9719	0.9726	0.9732	0.9738	0.9744	0.9750	0.9756	0.9761	0.9767
2.0	0.9772	0.9778	0.9783	0.9788	0.9793	0.9798	0.9803	0.9808	0.9812	0.9817
2.1	0.9821	0.9826	0.9830	0.9834	0.9838	0.9842	0.9846	0.9850	0.9854	0.9857
2.2	0.9861	0.9864	0.9868	0.9871	0.9875	0.9878	0.9881	0.9884	0.9887	0.9890
2.3	0.9893	0.9896	0.9898	0.9901	0.9904	0.9906	0.9909	0.9911	0.9913	0.9916
2.4	0.9918	0.9920	0.9922	0.9925	0.9927	0.9929	0.9931	0.9932	0.9934	0.9936
2.5	0.9938	0.9940	0.9941	0.9943	0.9945	0.9946	0.9948	0.9949	0.9951	0.9952
2.6	0.9953	0.9955	0.9956	0.9957	0.9959	0.9960	0.9961	0.9962	0.9963	0.9964
2.7	0.9965	0.9966	0.9967	0.9968	0.9969	0.9970	0.9971	0.9972	0.9973	0.9974
2.8	0.9974	0.9975	0.9976	0.9977	0.9977	0.9978	0.9979	0.9979	0.9980	0.9981
2.9	0.9981	0.9982	0.9982	0.9983	0.9984	0.9984	0.9985	0.9985	0.9986	0.9986
3.0	0.9987	0.9987	0.9987	0.9988	0.9988	0.9989	0.9989	0.9989	0.9990	0.9990
3.1	0.9990	0.9991	0.9991	0.9991	0.9992	0.9992	0.9992	0.9992	0.9993	0.9993
3.2	0.9993	0.9993	0.9994	0.9994	0.9994	0.9994	0.9994	0.9995	0.9995	0.9995
3.3	0.9995	0.9995	0.9995	0.9996	0.9996	0.9996	0.9996	0.9996	0.9996	0.9997
3.4	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9997	0.9998
3.5	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998	0.9998
3.6	0.9998	0.9998	0.9999							

TABLE A.2

t Distribution: Critical Values of t

<i>Degrees of freedom</i>	<i>Two-tailed test: One-tailed test:</i>	<i>Significance level</i>					
		10% 5%	5% 2.5%	2% 1%	1% 0.5%	0.2% 0.1%	0.1% 0.05%
1		6.314	12.706	31.821	63.657	318.309	636.619
2		2.920	4.303	6.965	9.925	22.327	31.599
3		2.353	3.182	4.541	5.841	10.215	12.924
4		2.132	2.776	3.747	4.604	7.173	8.610
5		2.015	2.571	3.365	4.032	5.893	6.869
6		1.943	2.447	3.143	3.707	5.208	5.959
7		1.894	2.365	2.998	3.499	4.785	5.408
8		1.860	2.306	2.896	3.355	4.501	5.041
9		1.833	2.262	2.821	3.250	4.297	4.781
10		1.812	2.228	2.764	3.169	4.144	4.587
11		1.796	2.201	2.718	3.106	4.025	4.437
12		1.782	2.179	2.681	3.055	3.930	4.318
13		1.771	2.160	2.650	3.012	3.852	4.221
14		1.761	2.145	2.624	2.977	3.787	4.140
15		1.753	2.131	2.602	2.947	3.733	4.073
16		1.746	2.120	2.583	2.921	3.686	4.015
17		1.740	2.110	2.567	2.898	3.646	3.965
18		1.734	2.101	2.552	2.878	3.610	3.922
19		1.729	2.093	2.539	2.861	3.579	3.883
20		1.725	2.086	2.528	2.845	3.552	3.850
21		1.721	2.080	2.518	2.831	3.527	3.819
22		1.717	2.074	2.508	2.819	3.505	3.792
23		1.714	2.069	2.500	2.807	3.485	3.768
24		1.711	2.064	2.492	2.797	3.467	3.745
25		1.708	2.060	2.485	2.787	3.450	3.725
26		1.706	2.056	2.479	2.779	3.435	3.707
27		1.703	2.052	2.473	2.771	3.421	3.690
28		1.701	2.048	2.467	2.763	3.408	3.674
29		1.699	2.045	2.462	2.756	3.396	3.659
30		1.697	2.042	2.457	2.750	3.385	3.646
32		1.694	2.037	2.449	2.738	3.365	3.622
34		1.691	2.032	2.441	2.728	3.348	3.601
36		1.688	2.028	2.434	2.719	3.333	3.582
38		1.686	2.024	2.429	2.712	3.319	3.566
40		1.684	2.021	2.423	2.704	3.307	3.551
42		1.682	2.018	2.418	2.698	3.296	3.538
44		1.680	2.015	2.414	2.692	3.286	3.526
46		1.679	2.013	2.410	2.687	3.277	3.515
48		1.677	2.011	2.407	2.682	3.269	3.505
50		1.676	2.009	2.403	2.678	3.261	3.496
60		1.671	2.000	2.390	2.660	3.232	3.460
70		1.667	1.994	2.381	2.648	3.211	3.435
80		1.664	1.990	2.374	2.639	3.195	3.416
90		1.662	1.987	2.368	2.632	3.183	3.402
100		1.660	1.984	2.364	2.626	3.174	3.390
120		1.658	1.980	2.358	2.617	3.160	3.373
150		1.655	1.976	2.351	2.609	3.145	3.357
200		1.653	1.972	2.345	2.601	3.131	3.340
300		1.650	1.968	2.339	2.592	3.118	3.323
400		1.649	1.966	2.336	2.588	3.111	3.315
500		1.648	1.965	2.334	2.586	3.107	3.310
600		1.647	1.964	2.333	2.584	3.104	3.307
∞		1.645	1.960	2.326	2.576	3.090	3.291

TABLE A.4

 χ^2 (Chi-Squared) Distribution: Critical Values of χ^2

<i>Degrees of freedom</i>	<i>Significance level</i>		
	5%	1%	0.1%
1	3.841	6.635	10.828
2	5.991	9.210	13.816
3	7.815	11.345	16.266
4	9.488	13.277	18.467
5	11.070	15.086	20.515
6	12.592	16.812	22.458
7	14.067	18.475	24.322
8	15.507	20.090	26.124
9	16.919	21.666	27.877
10	18.307	23.209	29.588