

# Assignment 2 for MAT4170

## Spline methods, Spring 2022

To be completed by Tuesday 1 March. Please send your solution as a single pdf file (including plots/figures) to `michaelf@math.uio.no`.

1. State the smoothness properties of  $B[0, 0, 0, 0, 1](x)$  and  $B[0, 1, 1, 1, 2](x)$ .
2. Use induction on the degree  $d$  to show that the formula

$$B[t_j, \dots, t_{j+d+1}](t_i) = B[t_j, \dots, t_{i-1}, t_{i+1}, \dots, t_{j+d+1}](t_i)$$

holds for  $i = j, \dots, j+d+1$ . Use it to prove property (5) of Lemma 2.3.

3. Do problem 2.17.
4. Do problem 2.18. Test your implementation on the knot vector

$$\mathbf{t} = (-1, -1, -1, -1, 0, 1, 2, 3, 4, 5, 5, 5, 5) \quad (1)$$

by plotting the non-zero B-splines of degree  $d = 3$  on the interval  $[0, 1]$ . Hint: do **not** use the B-spline matrices  $R_k(x)$ . Start with a vector  $B$  of zeros of length  $d + 1$  and set its last element to  $B_{\mu,0}(x) = 1$ . Then use the recurrence relation (2.1) to compute  $B_{j,k}(x)$  for  $j = \mu - k + 1, \dots, \mu$  for each  $k = 1, \dots, d$  (as in Figure 2.8), storing the result in  $B$ .

5. Do problem 2.19. Test your implementation by plotting the cubic spline with knot vector  $\mathbf{t}$  of (1) and coefficients

$$\mathbf{c} = (-1, 1, -1, 1, -1, 1, -1, 1, -1).$$

Hint: do **not** use the B-spline matrices  $R_k(x)$ . Start with the vector  $\mathbf{c} = (c_{\mu-d}, \dots, c_{\mu})$ . Then compute  $c_{j-k+1}$  by (2.20) for  $j = \mu, \dots, \mu - k + 1$  for each  $k = 1, \dots, d$  (as in Figure 2.9), storing the result in  $\mathbf{c}$ .

6. Do problem 2.26.