

MAT4520 - Spring 2024

Mandatory assignment 1 of 1

Submission deadline

Thursday 18th April 2024, 14:30 in Canvas (canvas.uio.no).

Instructions

Note that you have **one attempt** to pass the assignment. This means that there are no second attempts.

You can choose between scanning handwritten notes or typing the solution directly on a computer (for instance with \LaTeX). The assignment must be submitted as a single PDF file. Scanned pages must be clearly legible. The submission must contain your name, course and assignment number.

It is expected that you give a clear presentation with all necessary explanations. Remember to include all relevant plots and figures. All aids, including collaboration, are allowed, but the submission must be written by you and reflect your understanding of the subject. If we doubt that you have understood the content you have handed in, we may request that you give an oral account.

In exercises where you are asked to write a computer program, you need to hand in the code along with the rest of the assignment. It is important that the submitted program contains a trial run, so that it is easy to see the result of the code.

Application for postponed delivery

If you need to apply for a postponement of the submission deadline due to illness or other reasons, you have to contact the Student Administration at the Department of Mathematics (e-mail: studieinfo@math.uio.no) no later than the same day as the deadline.

All mandatory assignments in this course must be approved in the same semester, before you are allowed to take the final examination.

Complete guidelines about delivery of mandatory assignments:

uio.no/english/studies/admin/compulsory-activities/mn-math-mandatory.html

GOOD LUCK!

Problem 1.

In this problem we are going to construct a smooth manifold $G(k, n)$ - a so called Grassmannian - whose elements consists of the set of k -dimensional linear subspaces of \mathbb{R}^n . This is a generalization of a construction we have already considered, namely the construction of $\mathbb{R}\mathbb{P}^{n-1} = G(1, n)$.

A k -plane L in \mathbb{R}^n may be determined by choosing a basis a_1, \dots, a_k for L , and so L may be represented by an $n \times k$ -matrix

$$A = [a_1 a_2 \cdots a_k]$$

which we identify with an element of $\mathbb{R}^{n \cdot k}$. Since the vectors a_j are linearly independent we have that this matrix has rank k ¹. Letting $F(k, n)$ denote the set of $k \times n$ -matrices of rank k we have that any $B \in F(k, n)$ determines a k -plane. However, there is a large set of elements $B \in F(k, n)$ that determines the same k -plane. We introduce an equivalence relation \sim on $F(k, n)$ by declaring $A \sim B$ if and only if A and B determine the same k -plane. The manifold $G(k, n)$ will be constructed as the quotient $F(k, n)/\sim$ (where $F(k, n)$ has the topology induced by the identification with $\mathbb{R}^{n \cdot k}$).

- (i) Explain why $A \sim B$ if and only if there exists $g \in GL(k, \mathbb{R})$ such that $B = Ag$.
- (ii) Explain why for each fixed $g \in GL(k, \mathbb{R})$ the map $A \mapsto Ag$ is a homeomorphism on $F(k, n)$. Prove that \sim is an *open* equivalence relation.
- (iii) Prove that $G(k, n)$ is *second countable*.
- (iv) Prove that $G(k, n)$ is *Hausdorff*.

We would now like to find a smooth atlas for $G(k, n)$. For each multi-index I we let

$$V_I = \{A \in F(k, n) : \det(A_I) \neq 0\}$$

and note that $\{V_I\}$ is an open cover of $F(k, n)$. Since \sim is open we have that each $U_I = \pi(V_I)$ is an open subset of $G(k, n)$ and we have that $\{U_I\}$ is an open cover of $G(k, n)$.

Next we initially consider the fixed multi-index $I = (1, 2, \dots, k)$. For each $A \in V_I$ we have that

$$A \sim AA_I^{-1} = \begin{bmatrix} \text{Id} \\ A_I^{-1} \end{bmatrix}$$

¹If for a multi-index $I = (i_1, \dots, i_k), i_1 < i_2 < \cdots < i_k$, we let A_I denote the $k \times k$ -matrix whose rows are the I -rows of A , we have that the rank of A is k if and only if $\det A_I \neq 0$ for at least one I .

where $\widehat{I} = (k + 1, \dots, n)$. Therefore any equivalence class may be represented by an element of $\mathbb{R}^{(n-k) \times k}$.

(v) Show that the map

$$\tilde{\phi}_I : V_I \rightarrow \mathbb{R}^{(n-k) \times k}$$

induces a homeomorphism

$$\phi_I : U_I \rightarrow \mathbb{R}^{(n-k) \times k}$$

- (vi) Define similarly maps ϕ_I for all multi-indices I . Show that the transition maps $\phi_J \circ \phi_I^{-1}$ are smooth for multi-indices I, J .
- (vii) Conclude that we have given $G(k, n)$ the structure of a smooth manifold.