



UiO : Department of Mathematics
University of Oslo

STK4021

Examples

Ida Scheel

October 20, 2016

Overview of examples

1 Single-parameter models

- Epidemiology: Estimating a rate from Poisson data

2 Multi-parameter models

- Multinomial sampling distribution with a Dirichlet prior: Application to a US 2016 presidential election poll
- Analysis of a bioassay experiment

3 References

Single-parameter model for epidemiology

- Concerns estimating a rate from Poisson data (idealized example from the textbook [1], pp 45-46)
- Consider a survey of the causes of death in a single year for a city in the US
- Population 200.000, $y = 3$ persons died of asthma
 - Crude estimate of $3/200.000 = 1.5$ per 100.000 persons per year
- For epidemiological data like this, a Poisson sampling distribution is commonly used, assuming exchangeability given exposure and rate parameter
 - Let θ be the true, underlying long-term asthma mortality rate per 100.000 persons per year in the city
 - The exposure is $x = 2.0$ (since θ is defined per 100.000 persons per year)
 - Hence, the sampling distribution is $y \sim \text{Poisson}(2.0\theta)$

Prior distribution

- Asthma mortality rates around the world typically are around 0.6 per 100.000, and rarely above 1.5 per 100.000 in Western countries
- Assume exchangeability between this city and other Western cities, and this year and other years
- Know that Gamma(a, b) is the conjugate prior distribution, use that for convenience, must find suitable values of a and b that match the prior information
- Book: $\theta \sim \text{Gamma}(3.0, 5.0)$ (mean=0.6, 97.5% of the mass lies below 1.44, prior probability of $\theta < 1.5$ 98.0%)
- Slightly different (with more uncertainty) suggestion:
 $\theta \sim \text{Gamma}(1.2, 2.0)$ (mean=0.6, 97.5% of the mass lies below 2.05, prior probability of $\theta < 1.5$ 92.9%)
- “rarely above 1.5 per 100.000” is open for interpretation

Posterior distribution

- We know that the posterior distribution for θ will be $\text{Gamma}(a + y, b + x)$
- Book prior: Posterior is $\text{Gamma}(6.0, 7.0)$
- Alternative prior: Posterior is $\text{Gamma}(4.2, 4.0)$
- The two different priors yields somewhat different posterior distributions and conclusions (see R-script)
- Little data!!! Prior is influential

Posterior distribution with additional data

- Additional data: Suppose we now have 10 years of data, with 30 deaths caused by asthma over the 10 years. Assume the population size is constant at 200.000 over the period
- Now $y = 30$ and the exposure is $x = \frac{200.000 \times 10}{100.000} = 20$ (since θ is defined per 100.000 persons per year)
- Book prior: Posterior is Gamma(33.0, 25.0)
- Alternative prior: Posterior is Gamma(31.2, 22.0)
- The posterior results with the two different priors are more similar now with more data, but still slightly different (see R-script)
- More data, prior is less influential

Alternative specification with n independent outcomes

- Could alternatively say that $y_i, i = 1, \dots, n$ is the number of deaths caused by asthma per 100.000 persons per year
- Let θ still be the true, underlying long-term asthma mortality rate per 100.000 persons per year in the city
- Then

$$y_i \sim \text{Pois}(x_i\theta), i = 1, \dots, n$$

where the exposure is $x_i, i = 1, \dots, n$

- Then we know that the likelihood is

$$p(y | \theta) \propto \theta^{\sum_{i=1}^n y_i} e^{-\theta \sum_{i=1}^n x_i}$$

where in the example we have $y_i = 3, x_i = 2, i = 1, \dots, n$ for (i) $n = 1$ and (ii) $n = 10$

The normal approximation

- The normal approximation to the posterior distribution for θ based on $\log p(y | \theta) = C + \sum_{i=1}^n y_i \log \theta - \theta \sum_{i=1}^n x_i$ can easily be found. First find the mode

$$\frac{d \log p(y | \theta)}{d\theta} = \frac{\sum_{i=1}^n y_i}{\theta} - \sum_{i=1}^n x_i$$

which is =0 for $\theta = \hat{\theta} = \frac{\sum_{i=1}^n y_i}{\sum_{i=1}^n x_i}$

- Then the Fisher information (since we allow for different exposure-values x_i , the y_i 's are not iid):

$$\begin{aligned} n \cdot J(\theta) &= E \left[-\frac{d^2 \log p(y | \theta)}{d\theta^2} \mid \theta \right] = E \left[\frac{\sum_{i=1}^n y_i}{\theta^2} \right] \\ &= \frac{\sum_{i=1}^n x_i \theta}{\theta^2} = \frac{\sum_{i=1}^n x_i}{\theta} \end{aligned}$$

(using $E[y_i] = x_i \theta$)

The normal approximation

- Hence

$$\hat{\theta} = \frac{\sum_{i=1}^n y_i}{\sum_{i=1}^n x_i}$$
$$n \cdot J(\hat{\theta}) = \frac{(\sum_{i=1}^n x_i)^2}{\sum_{i=1}^n y_i}$$

and for large n

$$p(\theta | y) \approx N\left(\theta \mid \hat{\theta}, \left(n \cdot J(\hat{\theta})\right)^{-1}\right)$$

- For $n = 1$ the Normal approximation is $N\left(\frac{3}{2}, \frac{3}{2^2}\right) = N(1.5, 0.75)$
and for $n = 10$ it is $N\left(\frac{30}{20}, \frac{30}{20^2}\right) = N(1.5, 0.075)$

Multinomial sampling distribution with a Dirichlet prior

- Application: 2016 US presidential election poll (Sept-16)
 - $n = 911$ representative, likely voters were asked which candidate they prefer in the 2016 US presidential election
 - $y_1 = 392$ preferred Clinton, $y_2 = 364$ preferred Trump, and $y_3 = 155$ preferred other candidates or had no opinion
- Multinomial sampling distribution with
 - probability θ_1 of preferring Clinton
 - probability θ_2 of preferring Trump
 - probability θ_3 of preferring other candidates or having no opinion
 - $\sum_{i=1}^3 \theta_i = 1$ (hence there are in fact only two parameters)
- A non-informative uniform Dirichlet(1,1,1) prior for $(\theta_1, \theta_2, \theta_3)$
- Hence, the posterior distribution for $(\theta_1, \theta_2, \theta_3)$ is
Dirichlet($1 + y_1, 1 + y_2, 1 + y_3$) = Dirichlet(393, 365, 156)

The posterior distribution of an estimand of interest

- Suppose we are interested in the posterior distribution of $\frac{\theta_1}{\theta_2}$
- This can easily be approximated by for $i = 1, \dots, S$ doing
 - Sample $\theta^{(i)}$ from the posterior distribution of $(\theta_1, \theta_2, \theta_3)$
 - Compute $\frac{\theta_1^{(i)}}{\theta_2^{(i)}}$
- The S values of $\frac{\theta_1^{(i)}}{\theta_2^{(i)}}$ for $i = 1, \dots, S$ are then samples from the posterior distribution of $\frac{\theta_1}{\theta_2}$

The Normal approximation

- Our interest primarily lies in θ_1 and θ_2 , therefore we focus on these two parameters and replace θ_3 by $1 - \theta_1 - \theta_2$
- The likelihood for $\theta = (\theta_1, \theta_2)$ is

$$p(y | \theta) \propto \prod_{i=1}^3 \theta_i^{y_i} = \theta_1^{y_1} \cdot \theta_2^{y_2} \cdot (1 - \theta_1 - \theta_2)^{y_3}$$

- The Normal approximation to the posterior distribution for θ based on

$\log p(y | \theta) = C + y_1 \log \theta_1 + y_2 \log \theta_2 + y_3 \log (1 - \theta_1 - \theta_2)$ can easily be found. First find the mode

$$\frac{d \log p(y | \theta)}{d \theta_i} = \frac{y_i}{\theta_i} - \frac{y_3}{1 - \theta_1 - \theta_2}, i = 1, 2$$

which is =0 for $\theta_i = \hat{\theta}_i = \frac{y_i}{\sum_{j=1}^3 y_j}$

The Normal approximation

- Then the Fisher information:

$$\frac{d^2 \log p(y | \theta)}{d\theta_i^2} = -\frac{y_i}{\theta_i^2} - \frac{y_3}{(1 - \theta_1 - \theta_2)^2}, i = 1, 2$$

$$\frac{d^2 \log p(y | \theta)}{d\theta_1 d\theta_2} = -\frac{y_3}{(1 - \theta_1 - \theta_2)^2}$$

⇓

$$n \cdot J(\theta) = E \left[- \begin{pmatrix} -\frac{y_1}{\theta_1^2} - \frac{y_3}{(1-\theta_1-\theta_2)^2} & -\frac{y_3}{(1-\theta_1-\theta_2)^2} \\ -\frac{y_3}{(1-\theta_1-\theta_2)^2} & -\frac{y_2}{\theta_2^2} - \frac{y_3}{(1-\theta_1-\theta_2)^2} \end{pmatrix} \right]$$

$$= \begin{pmatrix} -\frac{n}{\theta_1} - \frac{n}{1-\theta_1-\theta_2} & -\frac{n}{1-\theta_1-\theta_2} \\ -\frac{n}{1-\theta_1-\theta_2} & -\frac{n}{\theta_2} - \frac{n}{1-\theta_1-\theta_2} \end{pmatrix}$$

(using $E[y_i] = n\theta_i$)

The Normal approximation

- Hence

$$\hat{\theta}_i = \frac{y_i}{\sum_{j=1}^3 y_j}$$
$$n \cdot J(\hat{\theta}) = n \cdot \begin{pmatrix} -\frac{1}{\hat{\theta}_1} & -\frac{1}{1-\hat{\theta}_1-\hat{\theta}_2} & -\frac{1}{1-\hat{\theta}_1-\hat{\theta}_2} \\ -\frac{1}{1-\hat{\theta}_1-\hat{\theta}_2} & \frac{1}{\hat{\theta}_2} & \frac{1}{1-\hat{\theta}_1-\hat{\theta}_2} \end{pmatrix}$$

and for large n

$$p(\theta | y) \approx N\left(\theta \mid \hat{\theta}, \left(n \cdot J(\hat{\theta})\right)^{-1}\right)$$

- Here we know the exact posterior distribution, can compare it to the Normal approximation by for example contour-plots (see R-script)

The application and sampling distribution

- Example from the textbook [1], section 3.7
- A bioassay experiment typically concerns giving various dose levels of a drug/chemical compound to a batch of animals and measure a binary response (alive/dead or tumor/no tumor)
- The data for k dose levels are of the form

$$(x_i, n_i, y_i), \quad i = 1, \dots, k$$

where x_i is the i 'th dose level given to n_i animals of which y_i animals responded with "success" (e.g. death)

- Reasonable to model the response of the animals within the i 'th group (given dose x_i) as exchangeable, by modelling them as independent with equal probabilities of success θ_i , i.e. a binomial model

$$y_i \mid \theta_i \sim \text{Bin}(n_i, \theta_i)$$

Logistic regression model for the probabilities

- The parameters $\theta_1, \dots, \theta_k$ should be not be modelled as exchangeable, since we have the dose levels x_1, \dots, x_k
- Rather model the pairs $\theta_i \mid x_i, i = 1, \dots, k$ by a logistic regression model

$$\text{logit}(\theta_i) = \alpha + \beta x_i, \quad i = 1, \dots, k$$

where $\text{logit}(\theta_i) = \log \frac{\theta_i}{1-\theta_i}$ is the logistic transformation

- Hence $\theta_i = \text{logit}^{-1}(\alpha + \beta x_i) = \frac{e^{\{\alpha + \beta x_i\}}}{1 + e^{\{\alpha + \beta x_i\}}}$ and the likelihood contribution from group i for the parameters α and β is

$$\begin{aligned} p(y_i \mid \alpha, \beta) &\propto \theta_i^{y_i} (1 - \theta_i)^{n_i - y_i} \\ &= \left(\frac{e^{\{\alpha + \beta x_i\}}}{1 + e^{\{\alpha + \beta x_i\}}} \right)^{y_i} \left(\frac{1}{1 + e^{\{\alpha + \beta x_i\}}} \right)^{n_i - y_i} \end{aligned}$$

Prior and posterior distributions

- We assume an improper prior distribution for the parameters α and β : $p(\alpha, \beta) \propto 1$
- Hence, α and β are independent a priori and marginally uniformly distributed
- Hence the joint posterior distribution for α and β can be expressed as

$$\begin{aligned} p(\alpha, \beta | y) &\propto p(\alpha, \beta) \prod_{i=1}^k p(y_i | \alpha, \beta, n_i, x_i) \\ &= \prod_{i=1}^k \left(\frac{e^{\{\alpha + \beta x_i\}}}{1 + e^{\{\alpha + \beta x_i\}}} \right)^{y_i} \left(\frac{1}{1 + e^{\{\alpha + \beta x_i\}}} \right)^{n_i - y_i} \end{aligned}$$

Data and graph of the model

Bioassay data from an experiment (table 3.1 from the textbook [1], see the textbook for reference)

Dose, x_i (log g/ml)	Number of animals n_i	Number of deaths y_i
-0.86	5	0
-0.30	5	1
-0.05	5	3
0.73	5	5




Figure: Graph representation of the model

Posterior analysis

- The normalised posterior distribution is not available analytically
- Hence, some numerical approximation must be performed, e.g. sampling
- Book, ch 3.7, compute the posterior density on a grid of points, then normalise by setting the total probability over the grid of points equal to 1
- Later we can do e.g. MCMC
- Now: Normal approximation (Exercise 2 in Chapter 4)

The normal approximation

- The ML-estimates of (α, β) can be found by using standard software for logistic regression, the results are (from the textbook) $(\hat{\alpha}, \hat{\beta}) = (0.8, 7.7)$
- Log-likelihood for one datapoint:

$$\begin{aligned}l_i &= \log p(y_i | \alpha, \beta) \\&= C + y_i \log \left(\frac{e^{\{\alpha + \beta x_i\}}}{1 + e^{\{\alpha + \beta x_i\}}} \right) + (n_i - y_i) \log \left(\frac{1}{1 + e^{\{\alpha + \beta x_i\}}} \right) \\&= C + y_i(\alpha + \beta x_i) - n_i \log \left(1 + e^{\{\alpha + \beta x_i\}} \right)\end{aligned}$$

- Hence

$$\begin{aligned}\frac{dl_i}{d\alpha} &= y_i - \frac{n_i e^{\{\alpha + \beta x_i\}}}{1 + e^{\{\alpha + \beta x_i\}}} \\ \frac{dl_i}{d\beta} &= y_i x_i - \frac{n_i x_i e^{\{\alpha + \beta x_i\}}}{1 + e^{\{\alpha + \beta x_i\}}}\end{aligned}$$

The normal approximation

- The second partial derivatives:

$$\begin{aligned}\frac{d^2 l_i}{d\alpha^2} &= - \frac{n_i e^{\{\alpha+\beta x_i\}} \left(1 + e^{\{\alpha+\beta x_i\}}\right) - n_i e^{\{\alpha+\beta x_i\}} e^{\{\alpha+\beta x_i\}}}{(1 + e^{\{\alpha+\beta x_i\}})^2} \\ &= - \frac{n_i e^{\{\alpha+\beta x_i\}}}{(1 + e^{\{\alpha+\beta x_i\}})^2}\end{aligned}$$

$$\begin{aligned}\frac{d^2 l_i}{d\beta^2} &= - \frac{n_i x_i^2 e^{\{\alpha+\beta x_i\}} \left(1 + e^{\{\alpha+\beta x_i\}}\right) - n_i x_i e^{\{\alpha+\beta x_i\}} x_i e^{\{\alpha+\beta x_i\}}}{(1 + e^{\{\alpha+\beta x_i\}})^2} \\ &= - \frac{n_i x_i^2 e^{\{\alpha+\beta x_i\}}}{(1 + e^{\{\alpha+\beta x_i\}})^2}\end{aligned}$$

$$\begin{aligned}\frac{d^2 l_i}{d\alpha d\beta} &= - \frac{n_i x_i e^{\{\alpha+\beta x_i\}} \left(1 + e^{\{\alpha+\beta x_i\}}\right) - n_i e^{\{\alpha+\beta x_i\}} x_i e^{\{\alpha+\beta x_i\}}}{(1 + e^{\{\alpha+\beta x_i\}})^2} \\ &= - \frac{n_i x_i e^{\{\alpha+\beta x_i\}}}{(1 + e^{\{\alpha+\beta x_i\}})^2}\end{aligned}$$

The normal approximation

- The normal approximation for (α, β) has mean $(\hat{\alpha}, \hat{\beta})$ and covariance matrix $(n \cdot J((\hat{\alpha}, \hat{\beta})))^{-1}$, where (remember that y_1, \dots, y_k are not identically distributed)

$$n \cdot J((\hat{\alpha}, \hat{\beta})) = \begin{pmatrix} \sum_{i=1}^k \frac{n_i e^{\{\hat{\alpha} + \hat{\beta} x_i\}}}{(1 + e^{\{\hat{\alpha} + \hat{\beta} x_i\}})^2} & \sum_{i=1}^k \frac{n_i x_i e^{\{\hat{\alpha} + \hat{\beta} x_i\}}}{(1 + e^{\{\hat{\alpha} + \hat{\beta} x_i\}})^2} \\ \sum_{i=1}^k \frac{n_i x_i e^{\{\hat{\alpha} + \hat{\beta} x_i\}}}{(1 + e^{\{\hat{\alpha} + \hat{\beta} x_i\}})^2} & \sum_{i=1}^k \frac{n_i x_i^2 e^{\{\hat{\alpha} + \hat{\beta} x_i\}}}{(1 + e^{\{\hat{\alpha} + \hat{\beta} x_i\}})^2} \end{pmatrix}$$

The normal approximation

- The normal approximation variances are the diagonal elements of $(n \cdot J((\hat{\alpha}, \hat{\beta})))^{-1}$, hence

$$\widehat{\text{Var}}(\alpha) = \frac{\sum_{i=1}^k \frac{n_i x_i^2 e^{\{\hat{\alpha} + \hat{\beta} x_i\}}}{(1 + e^{\{\hat{\alpha} + \hat{\beta} x_i\}})^2}}{\left(\sum_{i=1}^k \frac{n_i e^{\{\hat{\alpha} + \hat{\beta} x_i\}}}{(1 + e^{\{\hat{\alpha} + \hat{\beta} x_i\}})^2} \right) \left(\sum_{i=1}^k \frac{n_i x_i^2 e^{\{\hat{\alpha} + \hat{\beta} x_i\}}}{(1 + e^{\{\hat{\alpha} + \hat{\beta} x_i\}})^2} \right) - \left(\sum_{i=1}^k \frac{n_i x_i e^{\{\hat{\alpha} + \hat{\beta} x_i\}}}{(1 + e^{\{\hat{\alpha} + \hat{\beta} x_i\}})^2} \right)^2}$$

$$\widehat{\text{Var}}(\beta) = \frac{\sum_{i=1}^k \frac{n_i e^{\{\hat{\alpha} + \hat{\beta} x_i\}}}{(1 + e^{\{\hat{\alpha} + \hat{\beta} x_i\}})^2}}{\left(\sum_{i=1}^k \frac{n_i e^{\{\hat{\alpha} + \hat{\beta} x_i\}}}{(1 + e^{\{\hat{\alpha} + \hat{\beta} x_i\}})^2} \right) \left(\sum_{i=1}^k \frac{n_i x_i^2 e^{\{\hat{\alpha} + \hat{\beta} x_i\}}}{(1 + e^{\{\hat{\alpha} + \hat{\beta} x_i\}})^2} \right) - \left(\sum_{i=1}^k \frac{n_i x_i e^{\{\hat{\alpha} + \hat{\beta} x_i\}}}{(1 + e^{\{\hat{\alpha} + \hat{\beta} x_i\}})^2} \right)^2}$$

References I



A. Gelman, J. B. Carlin, H. Stern, D. B. Dunson, A. Vehtari and D. B. Rubin
Bayesian Data Analysis, Third edition.
Chapman&Hall/CRC Texts in statistical science, 2014.