

## Lecture note 3 (with corrections and references made 9 september 2013)

### General companion form, and the ADL derived from the VAR.

#### The companion form of a VAR

In Lecture 2 we saw that the VAR in two variables  $X_t$  and  $Y_t$ , and with first order dynamics, an VAR(1), is already on so called “companion form”. We now demonstrate that also a general VAR in  $n$  variables and dynamics of order  $p$  can be expressed as a VAR(1) with a companion matrix that contain all the parameters of the system.

We define  $\mathbf{y}_t$  as the  $n \times 1$  vector

$$\mathbf{y}_t = [Y_{1t}, Y_{2t}, \dots, Y_{nt}]'$$

and write the system with dynamics of degree  $p$  as:

$$\mathbf{y}_t = \phi_1 \mathbf{y}_{t-1} + \phi_2 \mathbf{y}_{t-2} + \dots + \phi_p \mathbf{y}_{t-p} + \boldsymbol{\epsilon}_t \quad (1)$$

where  $\phi_i$  is a  $n \times n$  matrix with parameters and  $\boldsymbol{\epsilon}_t$  is a vector of random variables (it can be jointly normally distributed (Gaussian) or at least stationary). The *companion form* of this general system is:

$$\underbrace{\begin{pmatrix} \mathbf{y}_t \\ \mathbf{y}_{t-1} \\ \vdots \\ \mathbf{y}_{t-p+1} \end{pmatrix}}_{\mathbf{z}_t \quad np \times 1} = \underbrace{\begin{pmatrix} \phi_1 & \phi_2 & \cdots & \phi_{p-1} & \phi_p \\ \mathbf{I} & \mathbf{0} & \mathbf{0} & \cdots & \mathbf{0} \\ \mathbf{0} & \mathbf{I} & \mathbf{0} & \cdots & \mathbf{0} \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ \mathbf{0} & \mathbf{0} & \mathbf{0} & \mathbf{I} & \mathbf{0} \end{pmatrix}}_{\mathbf{F}_{np \times np}} \underbrace{\begin{pmatrix} \mathbf{y}_{t-1} \\ \mathbf{y}_{t-2} \\ \vdots \\ \mathbf{y}_{t-p} \end{pmatrix}}_{\mathbf{z}_{t-1} \quad np \times 1} + \underbrace{\begin{pmatrix} \boldsymbol{\epsilon}_t \\ \mathbf{0} \\ \vdots \\ \mathbf{0} \end{pmatrix}}_{\boldsymbol{\epsilon}_t \quad np \times 1} \quad (2)$$

or, written compactly, with the suggested notation:

$$\mathbf{z}_t = \mathbf{F} \mathbf{z}_{t-1} + \boldsymbol{\epsilon}_t, \quad (3)$$

The vector variable  $\mathbf{z}_t$  has a global asymptotic stable solution and is stationary if all eigenvalues of  $\mathbf{F}$  from

$$|\mathbf{F} - \lambda \mathbf{I}| = 0 \quad (4)$$

are less than one in magnitude. This means all real roots must be between  $-1$  and  $1$ , and all complex roots must have modulus less than one.

#### The ADL model derived from a Gaussian VAR

The following system is an example of a first order Gaussian VAR in the two time series  $X_t$  and  $Y_t$ :

$$\begin{pmatrix} Y_t \\ X_t \end{pmatrix} = \begin{pmatrix} \pi_{10} \\ \pi_{20} \end{pmatrix} + \begin{pmatrix} \pi_{11} & \pi_{12} \\ \pi_{21} & \pi_{22} \end{pmatrix} \begin{pmatrix} Y_{t-1} \\ X_{t-1} \end{pmatrix} + \begin{pmatrix} \varepsilon_{yt} \\ \varepsilon_{xt} \end{pmatrix} \quad (5)$$

$$\begin{pmatrix} \varepsilon_{yt} \\ \varepsilon_{xt} \end{pmatrix} \sim N \left( \mathbf{0}, \begin{pmatrix} \sigma_x^2 & \sigma_{xy} \\ \sigma_{xy} & \sigma_y^2 \end{pmatrix} \mid Y_{t-1}, X_{t-1} \right). \quad (6)$$

The conditioning in (6) means that the normal distribution of  $\epsilon_t = (\varepsilon_{yt}, \varepsilon_{xt})'$  in the VAR is conditional on the history of the system up to period  $t - 1$ . In Lecture note 4, we generalize the argument by starting from the so called Haavelmo distribution, but as said in the lectures and computer class, it is enough to study the first order case carefully to get the right understanding.

Start by writing (5) as

$$Y_t = \mu_{y,t-1} + \varepsilon_{yt} \quad (7)$$

$$X_t = \mu_{x,t-1} + \varepsilon_{xt} \quad (8)$$

where the conditional expectations  $\mu_{y,t-1} \equiv E(Y_t | Y_{t-1}, X_{t-1})$  and  $\mu_{x,t-1} \equiv E(X_t | Y_{t-1}, X_{t-1})$  are given by

$$\mu_{y,t-1} = \pi_{10} + \pi_{11}Y_{t-1} + \pi_{12}X_{t-1} \quad (9)$$

$$\mu_{x,t-1} = \pi_{20} + \pi_{21}Y_{t-1} + \pi_{22}X_{t-1}. \quad (10)$$

$Y_t$  and  $X_t$  given by (7), (8) and (6) have a joint normal distribution which is conditional on  $X_{t-1}$  and  $Y_{t-1}$ . It follows from the properties of the normal distribution that the conditional distribution of  $Y_t$  given  $X_t$  is also normal, with expectation:

$$\begin{aligned} E(Y_t | X_t, X_{t-1}, Y_{t-1}) &= \mu_{y,t-1} - \rho_{xy} \frac{\sigma_y}{\sigma_x} \mu_{x,t-1} + \rho_{xy} \frac{\sigma_y}{\sigma_x} X_t \\ &= \pi_{10} - \frac{\omega_{xy}}{\sigma_x^2} \pi_{20} + \frac{\omega_{xy}}{\sigma_x^2} X_t + (\pi_{12} - \frac{\omega_{xy}}{\sigma_x^2} \pi_{22}) X_{t-1} \\ &\quad + (\pi_{11} - \frac{\omega_{xy}}{\sigma_x^2} \pi_{21}) Y_{t-1} \end{aligned}$$

where  $\rho_{xy}$  is the correlation coefficient between  $\varepsilon_{xt}$  and  $\varepsilon_{yt}$ :

$$\rho_{xy} = \frac{\sigma_{xy}}{\sqrt{\sigma_x^2} \sqrt{\sigma_y^2}}. \quad (11)$$

We can now define parameters:

$$\phi_0 = \pi_{10} - \frac{\sigma_{xy}}{\sigma_x^2} \pi_{20} \quad (12)$$

$$\phi_1 = \pi_{11} - \frac{\sigma_{xy}}{\sigma_x^2} \pi_{21} \quad (13)$$

$$\beta_0 = \frac{\sigma_{xy}}{\sigma_x^2} \quad (14)$$

$$\beta_1 = \pi_{12} - \frac{\sigma_{xy}}{\sigma_x^2} \pi_{22} \quad (15)$$

and write the conditional expectation as

$$E(Y_t | X_t, X_{t-1}, Y_{t-1}) = \phi_0 + \phi_1 Y_{t-1} + \beta_0 X_t + \beta_1 X_{t-1}. \quad (16)$$

Finally, define the disturbance  $\varepsilon_t$  as

$$\varepsilon_t = Y_t - E(Y_t | X_t, X_{t-1}, Y_{t-1}) \quad (17)$$

and write the conditional model for  $Y_t$  as

$$Y_t = \phi_0 + \phi_1 Y_{t-1} + \beta_0 X_t + \beta_1 X_{t-1} + \varepsilon_t \quad (18)$$

which is an ADL model, and the same as equation (13.58) in DM, but with the obvious change in notation (and with  $p = q = 1$ ).

Note that, again with reference to the normal distribution, the ADL-disturbance is:

$$\varepsilon_t = \varepsilon_{yt} - \frac{\sigma_{xy}}{\sigma_x^2} \varepsilon_{xt} \quad (19)$$

with variance:

$$\sigma^2 = \sigma_y^2(1 - \rho_{xy}^2). \quad (20)$$

If you guess that an ADL model with  $p$ -lags can be derived from a  $VAR(p)$  that condition on  $X_{t-p-1}$  and  $Y_{t-p-1}$  you are right! We loose nothing in generality about the status of the ADL as a conditional model by just looking at the simplest,  $VAR(1)$ , case. For those interested, we refer to *Lecture note 4*.

**Exogeneity and pre-determinedness:** Remember that the starting point is the joint distribution of  $\varepsilon_{yt}$  and  $\varepsilon_{xt}$  conditional on  $X_{t-1}$  and  $Y_{t-1}$ , cf (6). Therefore:

$$E(\varepsilon_t | X_{t-1}, Y_{t-1}) = 0$$

so the disturbance of the ADL model is uncorrelated with the conditioning variables of the VAR. But we have also

$$E(\varepsilon_t | \varepsilon_{xt}) = 0 \quad (21)$$

and since  $\varepsilon_{xt} = [X_t - E[X_t | X_{t-1}, Y_{t-1}]]$  by definition, (21) can alternatively be expressed as:

$$E(\varepsilon_t | X_t, X_{t-1}, Y_{t-1}) = 0 \quad (22)$$

showing that  $\varepsilon_t$  is uncorrelated with **all** the explanatory variables of the model, just like in the static regression models we have seen before. In sum:  $X_t, X_{t-1}, Y_{t-1}$  are exogenous variables, exactly in the sense given by (22).

However

$$E(\varepsilon_{t-j} | X_t, X_{t-1}, Y_{t-1}) \neq 0 \text{ for } j = 1, 2, \dots$$

since  $Y_{t-1}$  must depend on  $\varepsilon_{t-1}$  and earlier disturbances through the solution for  $Y_t$  obtained by repeated substitution of lagged  $Y_t$  in (18):

$$\begin{aligned} Y_t &= \phi_0 + \phi_1 Y_{t-1} + \beta_0 X_t + \beta_1 X_{t-1} + \varepsilon_t \\ &= \phi_0(1 + \phi_1) + X\text{-terms} + \varepsilon_t + \phi_1 \varepsilon_{t-1} + \dots + \phi_1^2 Y_{t-2} \end{aligned}$$

and so on. Hence  $Y_{t-1}$  is a pre-determined variable in (18), not an exogenous variable.

**Role of Gaussian VAR assumption:** In all important respects, the above remains valid if (6) is replaced by an IID assumption for the VAR disturbance. The only expectation is the equations that maps from the parameters of the normal distribution to the parameter of the ADL. But the parameters of the ADL will still be parameters in a conditional expectation (again, just as in the static/ordinary regression model case).